arXiv:hep-th/9301079v1 19 Jan 1993

November 1992
SUSX–TH–92/17.

Twisted Sector Yukawa Couplings For The $\mathbf{Z}_M \times \mathbf{Z}_N$ Orbifolds

D. Bailinλa, A. Loveλb and W.A. Sabra$\lambda c^{\star}$

λaSchool of Mathematical and Physical Sciences,
University of Sussex,
Brighton U.K.

λbDepartment of Physics,
Royal Holloway and Bedford New College,
University of London,
Egham, Surrey, U.K.

λcPhysics Department
Birkbeck College,
University of London,
Malet Street
London WC1E 7HX

ABSTRACT

The moduli dependent Yukawa couplings between twisted sectors of $\mathbf{Z}_M \times \mathbf{Z}_N$ Coxeter orbifolds are studied.

$\star$ Address from January 1993, Department of Physics, Royal Holloway and Bedford New College, University of London, Surrey, U.K.

1. Introduction

A knowledge of the Yukawa couplings for orbifold compactified string theory models [1, 2] is an important step towards comparison of the models with observation. Of particular importance is the exponential dependence of Yukawa couplings on moduli [3, 4] which can arise when all the states involved are in twisted sectors, because of its possible bearing on hierarchies [5]. Twisted sector Yukawa coulping have been investigated for the $\mathbf{Z}_3$ orbifold [3–6], for the $\mathbf{Z}_7$ orbifold [7] and for arbitrary $\mathbf{Z}_N$ orbifolds [8–11]. Here, we shall extend our recent discussion [12] of Yukawa couplings for the $\mathbf{Z}_3 \times \mathbf{Z}_3$ orbifold to all Coxeter $\mathbf{Z}_M \times \mathbf{Z}_N$ orbifolds with the point group realized by Coxeter elements acting on a direct sum of two-dimensional root lattices. After completion of the present work we have received a preprint by Stieberger [13] which also addresses the question of Yukawa couplings for $\mathbf{Z}_M \times \mathbf{Z}_N$ orbifolds though with the emphasis on general properties such as modular symmetries rather than specific orbifolds on specific lattices. A possible advantage that $\mathbf{Z}_M \times \mathbf{Z}_N$ orbifolds possess over $\mathbf{Z}_N$ orbifolds is that they can more generically provide the string loop threshold corrections needed to explain the low energy values of the gauge coupling constants [14–18] because they are not so tightly constrained by duality anomaly cancellation conditions [17].

2. Yukawa couplings for $\mathbf{Z}_M \times \mathbf{Z}_N$ Orbifolds.

To specify an orbifold model it is necessary to specify both the point group and the lattice on which the point group acts as a symmetry. Perhaps the most elegant possibility [19] is that the point group acts as a Coxeter element (product of Weyl reflections) on a root lattice of a Lie algebra. Here, we shall study the case where the root lattice involved is a direct sum of three 2-dimensional root lattices. For the $\mathbf{Z}_M \times \mathbf{Z}_N$ orbifolds (table 1) the factors involved are $\mathbf{Z}_3$, $\mathbf{Z}_4$, $\mathbf{Z}_6$ and $\mathbf{Z}_2$. These discrete groups can be realised as the Coxeter elements for the $SU(3)$, $SO(5)$, G_2 and $SO(4)$ root lattices, respectively [20, 19]. The $\mathbf{Z}_2$ group can also be realized as the square of the Coxeter element of the $SO(5)$ lattice and the cube of the Coxeter element on the G_2 lattice, and the $\mathbf{Z}_3$ group can also be realized as the square of the Coxeter element on the G_2 lattice. The simplest possibility (table 1 of the present work, and table 1 of ref. 20) is to realize $\mathbf{Z}_3$, $\mathbf{Z}_4$ and $\mathbf{Z}_6$ as a Coxeter element on each of the two dimensional sub-lattices (rather than as a power of Coxeter element). This can be done for each of the $\mathbf{Z}_M \times \mathbf{Z}_N$ orbifolds, except $\mathbf{Z}_3 \times \mathbf{Z}_6$, by the choice of lattices in table 1, and for $\mathbf{Z}_3 \times \mathbf{Z}_6$ it can be done for 2 of the 3 sub-lattices with $\mathbf{Z}_3$ realized as the square of the Coxeter element on the third G_2 sub-lattice. These are the $\mathbf{Z}_M \times \mathbf{Z}_N$ models we shall study here.

The action of the Coxeter element $C(G)$ of a rank 2 Lie algebra G on the basis vectors e_1 and e_2 of the root lattices may be taken to be

$$C(SU(3))e_1 = e_2, \quad C(SU(3))e_2 = -e_1 - e_2, \tag{2.1}$$

$$C(SO(5))e_1 = e_1 + 2e_2, \quad C(SO(5))e_2 = -e_1 - e_2, \tag{2.2}$$

$$C(G_2)e_1 = -e_1 - e_2, \quad C(G_2)e_2 = 3e_1 + 2e_2, \tag{2.3}$$

and

$$C(SO(4))e_1 = -e_1, \quad C(SO(4))e_2 = -e_2 \tag{2.4}$$

realizing $\mathbf{Z}_3$, $\mathbf{Z}_4$, $\mathbf{Z}_6$ and $\mathbf{Z}_2$, respectively. The powers of Coxeter elements $C\lambda 2(SO(5))$ and $C\lambda 3(G_2)$ realizing $\mathbf{Z}_2$ have the same action as $C(SO(4))$, and $C\lambda 2(G_2)$ realizing $\mathbf{Z}_3$ has the action

$$C\lambda 2(G_2)e_1 = -2e_1 - e_2, \quad C\lambda 2(G_2)e_2 = 3e_1 + e_2. \tag{2.5}$$

The non-zero Yukawa couplings obey point group selection rules, and selection rules for the H-lattice momentum associated with bosonized right-moving NSR fermionic degrees of freedom. These selection rules have already been written down [20] for $\mathbf{Z}_M \times \mathbf{Z}_N$ orbifolds. They also obey space group selection rules [3,4] which may be written in the following form. Let (α, l_1), (β, l_2) and (γ, l_3) be space group elements associated with the α, β and γ twisted sectors, where

$$l_1 = (I - \alpha) f_\alpha + (I - \alpha)\Lambda, \tag{2.6}$$

$$l_2 = (I - \beta) f_\beta + (I - \beta)\Lambda, \tag{2.7}$$

and

$$l_3 = (I - \gamma) f_\gamma + (I - \gamma)\Lambda. \tag{2.8}$$

In (2.7)–(2.8), f_α, f_β and f_γ are fixed points or fixed tori for the α, β and γ twisted sectors, respectively, and in each case Λ denotes an arbitrary lattice vector. Then, given that we have already satisfied the point group selection rule

$$\alpha\beta\gamma = I, \tag{2.9}$$

the full space group selection rule contains the additional condition

$$l_1 + l_2 + l_3 \text{ contains } 0. \tag{2.10}$$

The values of the non-zero Yukawa couplings amongst the twisted sectors are determined in detail by three–point functions involving fermionic and bosonic degrees of freedom. However, the crucial dependence on the deformation parameters (moduli) and the particular fixed points and fixed tori is entirely contained in (bosonic) twist field correlation functions [3,4] of the type

$$\mathcal{Z} = \prod_{i=1} \lambda 3 < \sigma_\alpha \lambda i(z_1, \bar{z}_1) \sigma_\beta \lambda i(z_2, \bar{z}_2) \sigma_\gamma \lambda i(z_3, \bar{z}_3) >= \prod_{i=1} \lambda 3 \mathcal{Z}_i, \tag{2.11}$$

where α, β and γ now label the twisted sectors and also the fixed points or fixed tori involved, and the index i distinguishes the twist fields associated with the complex coordinates X_i, $i = 1, 2, 3$, that define the six-dimensional compact manifold. The correlation function

$$\mathcal{Z}_i =< \sigma_\alpha \lambda i(z_1, \bar{z}_1)\sigma_\beta \lambda i(z_2, \bar{z}_2)\sigma_\gamma \lambda i(z_3, \bar{z}_3) > \tag{2.12}$$

factors [3, 4] into a quantum piece $\mathcal{Z}\lambda i_{qu}$ and a classical piece with all the dependence on the moduli and the particular fixed points and fixed tori involved contained in the classical piece.

$$\mathcal{Z}_i = \mathcal{Z}\lambda i_{qu} \sum_{X_{cl}} e\lambda{-}S\lambda i_{cl}, \tag{2.13}$$

where the classical action is

$$S\lambda i_{cl} = \frac{1}{\pi}\int d\lambda 2z\Big(\frac{\partial X_i}{\partial z}\frac{\partial \bar{X}_i}{\partial \bar{z}} + \frac{\partial X_i}{\partial \bar{z}}\frac{\partial \bar{X}_i}{\partial z}\Big). \tag{2.14}$$

Because of the string equations of motion

$$\frac{\partial\lambda 2X_i}{\partial z\partial \bar{z}} = 0, \tag{2.15}$$

$\partial X_i/\partial z$ and $\partial X_i/\partial \bar{z}$ are functions of z and $\bar{z}$ alone, respectively, which have to be chosen to respect the boundary conditions at z_1, z_2 and z_3 implicit in the operator product expansions with the twist fields [3]. Let the space group elements for the fixed points or fixed tori involved in the Yukawa coupling be (α, l_1), (β, l_2) and (γ, l_3), as before, with l_1, l_2 and l_3 as in (2.6)–(2.8). Also let P be the least common denominator for the fractional twists involved so that the action of the point group elements α, β and γ on the three complex coordinates X_i may be written in the form

$$\begin{aligned} \alpha &: X_i \to e\lambda 2\pi i A_i/P X_i, \\ \beta &: X_i \to e\lambda 2\pi i B_i/P X_i, \\ \gamma &: X_i \to e\lambda 2\pi i C_i/P X_i, \end{aligned} \tag{2.16}$$

where A_i, B_i and C_i are positive integers smaller than P with

$$A_i + B_i + C_i = 0 \ (\mathrm{mod}\ P), \tag{2.17}$$

for consistency with the point group selection rule. When one of the A_i, B_i or C_i is zero, that is one of the α, β or γ leaves the i-th complex plane invariant, then the three twist field correlation function reduces to a 2 twist field correlation function, which may be normalized to one. Otherwise, we need to evaluate the classical action using

$$\frac{\partial X_i}{\partial z} = a_i(z-z_1)\lambda-(1-A_i/P)(z-z_2)\lambda-(1-B_i/P)(z-z_3)\lambda-(1-C_i/P) \tag{2.18}$$

and

$$\frac{\partial X_i}{\partial \bar{z}} = b_i(\bar{z}-\bar{z}_1)\lambda-A_i/P(\bar{z}-\bar{z}_2)\lambda-B_i/P(\bar{z}-\bar{z}_3)\lambda-C_i/P. \tag{2.19}$$

In all the cases we shall study here, only the holomorphic field $\partial X_i/\partial z$ is an acceptable classical solution, because $\partial X_i/\partial \bar{z}$ gives a divergent contribution to the classical action and we must set

$$b_i = 0. \tag{2.20}$$

The constants a_i may be evaluated by the use of global monodromy conditions [3] where we integrate around a closed contour $\mathcal{C}_i$ such that X_i is shifted by v_i but not rotated. Thus,

$$\oint_{\mathcal{C}_i} dz \frac{\partial X_i}{\partial z} = v_i, \tag{2.21}$$

where we have dropped the contribution of $\partial X_i/\partial \bar{z}$ as just discussed. If, for example, we choose $\mathcal{C}_i$ to go B_i times around z_1 counter-clockwise and A_i times around z_2 clockwise, with the choice (allowed by $SL(2,C)$ symmetry)

$$z_1 = 0, \quad z_2 = 1, \quad z_3 = \infty, \tag{2.22}$$

then X_i is not rotated. Multiplying together the relevant space group elements, $(\alpha, l_1)\lambda B_i(\beta\lambda-1, l_2)\lambda A_i$ we find the shift in i-th complex plane to be the projection into this plane of

$$W_i = (1 - \alpha\lambda B_i)(f_\alpha - f_\beta + \Lambda), \tag{2.23}$$

where Λ is an arbitrary lattice vector. In the first instance, (2.23) is in the lattice basis for the Coxeter orbifold. If we transform to the orthonormal basis in which the action of α, β and γ is given by (2.16), then v_i is the component of W_i in the i-th complex direction, e.g. for X_1 the shift v_1 is the component of W_1 in the $1+i2$ direction, where 1 and 2 denote the orthonormal basis. The global monodromy condition may be evaluated with the aid of the integral

$$\oint_{\mathcal{C}_i} dz z\lambda-(1 - A_i y)(z-1)\lambda-(1 - B_i y) = -2i\sin(A_i B_i \pi y)\frac{\Gamma(A_i y)\Gamma(B_i y)}{\Gamma(A_i y + B_i y)} \tag{2.24}$$

with the result

$$2ia_i \sin(\frac{A_i B_i \pi}{P})\frac{\Gamma(\frac{A_i}{P})\Gamma(\frac{B_i}{P})}{\Gamma(\frac{A_i}{P} + \frac{B_i}{P})}(-z_\infty)\lambda-(1 - C_i/P) = v_i \tag{2.25}$$

which determines a_i. The contribution to the classical action may now be obtained by using Appendix A of [21], with the result

$$S_{cl}\lambda i = \frac{|v_i|\lambda 2}{4\pi \sin\lambda 2(\frac{A_i B_i \pi}{P})}\frac{\sin(\frac{A_i\pi}{P})\sin(\frac{B_i\pi}{P})}{\sin(\frac{(A_i + B_i)\pi}{P})}. \tag{2.26}$$

Exactly similar expressions may be written down when the contour is chosen instead to encircle the pair of twist fields associated with (α, l_1) and (γ, l_3), or the pair of twist fields associated with (β, l_2) and (γ, l_3). In general, to obtain consistency between these different global monodromy conditions it is necessary in (2.13) to restrict the sum over the initially arbitrary lattice vectors arising in v_i in a way which depends on the fixed tori involved. This problem has been solved for the case of $\mathbf{Z}_N$ orbifolds in appendix B of the last reference of [11], and this same result can be applied here provided we restrict attention to one particular complex coordinate X_i at a time.

In general, after general GSO projections inclusive of Wilson lines [22, 2] the surviving physical states are eigenstates of the two point group elements θ and ω generating the $\mathbf{Z}_M \times \mathbf{Z}_N$ point group. In the construction of these θ and ω eigenstates account must be taken of the fact that fixed points (or tori) are not always left invariant [23, 9] by the action of θ or ω. If m and n are the least integers such that for the fixed point (or torus) f_{kl} of $\theta\lambda k\omega\lambda l$,

$$\theta\lambda m f_{kl} \sim f_{kl} \tag{2.27}$$

and

$$\omega\lambda n f_{kl} \sim f_{kl}, \tag{2.28}$$

where $\sim$ signifies up to a lattice vector, then the θ and ω eigenstate combinations of fixed points $|p, q>$ are of the form

$$|p, q >= \sum_{r=0} \lambda m - 1 \sum_{s=0} \lambda n - 1 e\lambda{-i2\pi pr/m}e\lambda{-i2\pi qs/n}|\theta\lambda r\omega\lambda s f_{kl} >, \quad p = 0, .., m-1 \quad q = 0, .., n-1, \tag{2.29}$$

which are simultaneous eigenstates of θ and ω with eigenvalues $e\lambda i2\pi p/m$ and $e\lambda i2\pi q/n$, respectively.

Then, the Yukawa couplings for physical states are obtained as combinations of terms for definite fixed tori, subject to the space group selection rules.

3. The $\mathbf{Z}_2 \times \mathbf{Z}_6$ orbifold

In this section, we shall illustrate the calculations and results by considering the case of the $\mathbf{Z}_2 \times \mathbf{Z}_6$ Coxeter orbifold on the $SO(4) \times G_2\lambda 2$ lattice as in table 1. The corresponding results for the other $\mathbf{Z}_M \times \mathbf{Z}_N$ orbifolds are given in summary in the appendices. In the orthonormal basis, the action of the generators θ and ω of the point group on the complex string degrees of freedom X_i, $i = 1, 2, 3$, is

$$\theta X_1 = -X_1, \quad \theta X_2 = X_2, \quad \theta X_3 = -X_3, \tag{3.1}$$

and

$$\omega X_1 = X_1, \quad \omega X_2 = e\lambda 2\pi i/6X_2, \quad \omega X_3 = e\lambda{-}2\pi i/6X_3. \tag{3.2}$$

On the other hand, for this Coxeter orbifold, the action of θ and ω on the lattice basis is given by

$$\theta = (C(SO(4)), I, C\lambda{-}3(G_2)) \tag{3.3}$$

and

$$\omega = (I, C(G_2), C\lambda{-}1(G_2)), \tag{3.4}$$

where $C(H)$ is the Coxeter element for the Lie algebra of H. Explicitly, using (2.3) and (2.4), the action on the 6 basis vectors e_1,...,e_6 for the lattice basis is

$$\begin{array}{l} \theta e_1 = -e_1, \\ \theta e_2 = -e_2 \end{array}, \quad \begin{array}{l} \theta e_3 = e_3, \\ \theta e_4 = e_4 \end{array}, \quad \begin{array}{l} \theta e_5 = -e_5, \\ \theta e_6 = -e_6 \end{array} \tag{3.5}$$

and

$$\begin{array}{l} \omega e_1 = e_1, \\ \omega e_2 = e_2 \end{array}, \quad \begin{array}{l} \omega e_3 = -e_3 - e_4, \\ \omega e_4 = 3e_3 + 2e_4 \end{array}, \quad \begin{array}{l} \omega e_5 = 2e_5 + e_6, \\ \omega e_6 = -3e_5 - e_6. \end{array} \tag{3.6}$$

More generally, the original rigid lattice may be deformed in ways which preserve the action (3.5) and (3.6) of the point group. To obtain the most general choice of lattice compatible with the point group we must require that all the scalar products $e_i.e_j$ are preserved by the transformations (3.5) and (3.6). If we write

$$e_i.e_j = |e_i||e_j|\cos\theta_{ij}, \quad i \neq j, \tag{3.7}$$

we find that

$$|e_4| = \sqrt{3}|e_3| \equiv \sqrt{3}R_3, \tag{3.8}$$

$$|e_6| = \sqrt{3}|e_5| \equiv \sqrt{3}R_5, \tag{3.9}$$

and

$$\cos\theta_{34} = \cos\theta_{56} = -\frac{\sqrt{3}}{2}, \tag{3.10}$$

with all other $\cos\theta_{ij}$ zero except $\cos\theta_{12}$, and

$$|e_1| \equiv R_1, \tag{3.11}$$

$$|e_2| \equiv R_2, \tag{3.12}$$

and $\cos\theta_{12}$ undetermined. Thus we may take 5 independent deformations of the lattice compatible with the point group to be R_1, R_2, R_3, R_5 and $\cos\theta_{12}$. These are the deformation parameters or moduli. The fixed points and fixed tori for the twisted sectors are readily obtained using the action (3.5) and (3.6) of the generators of the point group on the (deformed) lattice. These are presented in table 2, for the twisted sectors containing massless states together with the associated space group elements. The space group selection rules for the various Yukawa couplings amongst twisted sectors are then easily obtained from (2.10). All Yukawa couplings consistent with the point group selection rule and H-momentum conservation rule for the bosonized NSR fermion right movers have already been listed in [20]. As already discussed in section 2, the physical states surviving generalized GSO projections are θ and ω eigenstates, which particular eigenstates survive depending on the details of the point group embeddings and Wilson lines in a particular model. The θ and ω eigenstate combinations of fixed points (or tori) may be obtained from the action of θ and ω on the fixed tori given in table 3. An orthonormal basis $\tilde{e}_1, ..., \tilde{e}_6$ in which X_1, X_2 and X_3 are the components of the string degrees of freedom in the $1+i2$, $2+i3$ and $5+i6$ directions, respectively, is obtained by comparing (3.1) and (3.2) with (3.5) and (3.6), and using (3.8)–(3.10) when ensuring the orthonormality of the basis. A suitable choice is related to the lattice basis by

$$\begin{array}{lll} e_1 = R_1\tilde{e}_1, & e_3 = R_3\tilde{e}_3, & e_5 = R_5\tilde{e}_5, \\ e_2 = R_2(\cos\theta_{12}\tilde{e}_1 + \sin\theta_{12}\tilde{e}_2), & e_4 = -R_3(\frac{3}{2}\tilde{e}_3 + \frac{\sqrt{3}}{2}\tilde{e}_4), & e_6 = -R_5(\frac{3}{2}\tilde{e}_5 + \frac{\sqrt{3}}{2}\tilde{e}_6). \end{array} \tag{3.13}$$

The moduli and fixed point (or tori) dependence of the non-zero Yukawa couplings for twisted sectors containing massless states may now be calculated using the approach described in section 2. Four types of contribution to the Yukawa coupling arise for the $\mathbf{Z}_2 \times \mathbf{Z}_6$ orbifold and also for the $\mathbf{Z}_3 \times \mathbf{Z}_6$, $\mathbf{Z}_2 \times \mathbf{Z}_6'$ and $\mathbf{Z}_6 \times \mathbf{Z}_6$ orbifolds. First, if $\mathcal{Z}_i$ of (2.12) reduces to a two point function because the i-th complex plane is unrotated in one of the three twisted sectors then we may normalize $\mathcal{Z}_i$ to 1. On the other hand, if all the three twist fields in (2.12) are non-trivial then the contribution to $S\lambda i_{cl}$ takes one of three forms. When all three twists are $e\lambda 2\pi i/3$, then

$$S\lambda i_{cl} = \frac{R\lambda 2_{2i-1}}{8\pi\sqrt{3}}\Big[(2m_{2i-1} - 2n_{2i-1} + 3k_{2i-1} + 6k_{2i})\lambda 2 + 3k_{2i-1}\lambda 2\Big] \tag{3.14}$$

where $i = 1, 2$ or 3, the arbitrary integers k_{2i-1} and k_{2i} derive from the freedom in the choice of the space group elements associated with particular fixed tori, and the integers m_{2i-1} and n_{2i-1} characterize any two of the fixed tori involved. The leading contribution to the Yukawa coupling in (2.13) is obtained by minimizing $S\lambda i_{cl}$ by varying the arbitrary integers k_{2i-1} and k_{2i}. In the case of (3.14), the minimum value of $S\lambda i_{cl}$ is given by

$$\begin{aligned}\left(S_{cl}\lambda i\right)_{min} =&0, \qquad m_{2i-1}-n_{2i-1}=0\\ =&\frac{2R_{2i-1}\lambda 2}{4\pi\sqrt{3}}, \qquad m_{2i-1}-n_{2i-1}=\pm 1,\pm 2.\end{aligned} \tag{3.15}$$

The corresponding moduli dependent suppression factor in the Yukawa coupling is $\exp\left(-S\lambda i_{cl}\right)_{min}$.

For one twist of $e\lambda 4\pi i/3$ and two twists of $e\lambda 2\pi i/6$,

$$S\lambda i_{cl}=\frac{R\lambda 2_{2i-1}}{16\pi\sqrt{3}}\left[(2n_{2i-1}+3k_{2i-1}+6k_{2i})\lambda 2+3k_{2i-1}\lambda 2\right] \tag{3.16}$$

where n_{2i-1} characterizes the fixed torus in the sector where the twist on X_i is $e\lambda 4\pi i/3$. In this case, the minimum value of $S\lambda i_{cl}$ is given by

$$\begin{aligned}\left(S_{cl}\lambda i\right)_{min} =&0, \qquad n_{2i-1}=0\\ =&\frac{R_{2i-1}\lambda 2}{4\pi\sqrt{3}}, \qquad n_{2i-1}=\pm 1.\end{aligned} \tag{3.17}$$

Finally, for twists of $e\lambda 2\pi i/6$, $e\lambda 2\pi i/3$ and -1,

$$S\lambda i_{cl}=\frac{R\lambda 2_{2i-1}}{16\pi\sqrt{3}}\left[(2n_{2i-1}+3p_{2i}+6k_{2i})\lambda 2+3(2n_{2i-1}+3p_{2i}-2p_{2i-1}+6k_{2i}+4k_{2i-1})\lambda 2\right], \tag{3.18}$$

where n_{2i-1} characterizes the fixed torus in the sector where the twist on X_i is $e\lambda 2\pi i/3$, and p_{2i-1} and p_{2i} characterize the fixed torus in the sector where the twist is -1. In this case, the minimum value of $S\lambda i_{cl}$ is given by

$$\begin{aligned}\left(S_{cl}\lambda i\right)_{min} =&0, \qquad n_{2i-1}=p_{2i-1}=p_{2i}=0\\ =&\frac{3R_{2i-1}\lambda 2}{4\pi\sqrt{3}}, \qquad n_{2i-1}=0,\ p_{2i-1},p_{2i}\text{ not both zero}\\ =&\frac{4R_{2i-1}\lambda 2}{4\pi\sqrt{3}}, \qquad n_{2i-1}=\pm 1,\ p_{2i-1}=p_{2i}=0\\ =&\frac{R_{2i-1}\lambda 2}{4\pi\sqrt{3}}, \qquad n_{2i-1}=\pm 1,\ p_{2i-1},p_{2i}\text{ not both zero.}\end{aligned} \tag{3.19}$$

The suppression factors for the non-zero Yukawa couplings amongst the twisted sectors of the $\mathbf{Z}_2 \times \mathbf{Z}_6$ orbifold are tabulated in table 4. Corresponding results for the other $\mathbf{Z}_M \times \mathbf{Z}_N$ orbifolds are given in the appendix. For the $\mathbf{Z}_2 \times \mathbf{Z}_4$ and $\mathbf{Z}_4 \times \mathbf{Z}_4$ orbifolds, the only non-trivial suppression factor arises when the twists on X_i are $e\lambda 2\pi i/4$, $e\lambda 2\pi i/4$ and -1. Then,

$$S\lambda i_{cl} = \frac{R\lambda 2_{2i-1}}{16\pi}\Big[(2n_{2i}+2p_{2i-1}-p_{2i}+4k_{2i-1}+2k_{2i})\lambda 2+(p_{2i}-2k_{2i})\lambda 2\Big], \tag{3.20}$$

where n_{2i} characterizes the fixed torus in one of the sectors where the twist on X_i is $e\lambda 2\pi i/4$ and p_{2i-1} and p_{2i} characterize the fixed torus in the sector where the twist is -1. In this case, the minimum value of $S\lambda i_{cl}$ is given by

$$\begin{aligned}\Big(S_{cl}\lambda i\Big)_{min} =& \frac{R_{2i-1}\lambda 2}{8\pi}, \qquad p_{2i}=1\\ =&0, \qquad p_{2i}=0,\ n_{2i}+p_{2i-1}=0\ (\mathrm{mod}\ 2)\\ =&\frac{R_{2i-1}\lambda 2}{4\pi}, \qquad p_{2i}=0,\ n_{2i}+p_{2i-1}=1\ (\mathrm{mod}\ 2).\end{aligned} \tag{3.21}$$

For all cases other than the $\mathbf{Z}_2 \times \mathbf{Z}_4$ and $\mathbf{Z}_4 \times \mathbf{Z}_4$ orbifolds, the suppression factors are written in the form $e\lambda-(\lambda_{2i-1}R_{2i-1}\lambda 2/4\pi\sqrt{3})$, and for the $\mathbf{Z}_2 \times \mathbf{Z}_4$ and $\mathbf{Z}_4 \times \mathbf{Z}_4$ orbifolds in the form $e\lambda-(\lambda_{2i-1}R_{2i-1}\lambda 2/16\pi)$. The suppression factors arising from (3.19) are summarized by

$$\lambda_i = \mu(n_{2i-1}, p_{2i-1}, p_{2i}), \tag{3.22}$$

where

$$\begin{aligned}\mu(n_{2i-1}, p_{2i-1}p_{2i}) =&0, \qquad n_{2i-1}=p_{2i-1}=p_{2i}=0\\ =&3, \qquad n_{2i-1}=0,\ p_{2i-1},p_{2i} \text{ not both zero}\\ =&4, \qquad n_{2i-1}=\pm 1,\ p_{2i-1}=p_{2i}=0\\ =&1, \qquad n_{2i-1}=\pm 1,\ p_{2i-1},p_{2i} \text{ not both zero,}\end{aligned} \tag{3.23}$$

where n_{2i-1} characterizes the fixed torus in the sector where the twist on X_i is $e\lambda 2\pi i/3$, and p_{2i-1} and p_{2i} characterize the fixed torus in the sector where the twist is -1. The suppression factors arising from (3.21) are summarized by

$$\lambda_i = \rho(n_{2i}, p_{2i-1}, p_{2i}), \tag{3.24}$$

where

$$\begin{aligned} \rho(n_{2i}, p_{2i-1}, p_{2i}) =& 2, \qquad p_{2i} = 1 \\ =& 0, \qquad p_{2i} = 0,\ n_{2i} + p_{2i-1} = 0 \ (\mathrm{mod}\ 2) \\ =& 4, \qquad p_{2i} = 0,\ n_{2i} + p_{2i-1} = 1 \ (\mathrm{mod}\ 2). \end{aligned} \tag{3.25}$$

The overall moduli and fixed tori independent normalization is easily obtained from the formula derived for $\mathbf{Z}_N$ orbifolds in terms of twists in the two-dimensional subspaces [11] which also applies here.

4. Conclusions

The moduli dependence of the Yukawa couplings amongst twisted sectors has been investigated for all $\mathbf{Z}_M \times \mathbf{Z}_N$ orbifolds with the point group realized in the simplest way by Coxeter elements acting on a direct sum of two-dimensional root lattices. All Yukawa couplings have a moduli dependence of the general form (3.14), (3.16), (3.18) or (3.20). A tabulation has been made of the moduli dependent suppression factors for all the non-zero Yukawa couplings. For any definite model, with specified point group embeddings and Wilson lines, the results of this paper may be employed to study suppression factors in the quark and lepton mass matrices.

ACKNOWLEDGEMENTS

This work was supported in part by S.E.R.C. We would like to thank S. Stieberger for a helpful communication.

REFERENCES

1. L. Dixon, J. A. Harvey, C. Vafa and E. Witten, Nucl. Phys. B261 (1985) 678; B274 (1986) 285.
2. A. Font, L. E. Ibanez, F. Quevedo and A. Sierra, Nucl. Phys. B331 (1991) 421.
3. L. Dixon, D. Friedan, E. Martinec and S. Shenker, Nucl. Phys. B282 (1987) 13.
4. S. Hamidi, and C. Vafa, Nucl. Phys. B279 (1987) 465.
5. L. E. Ibanez, Phys. Lett. B181 (1986) 269.
6. J. A. Casas and C. Munoz, Nucl. Phys. B332 (1990) 189
7. J. A. Casas, F. Gomez and C. Munoz, Phys. Lett. B251(1990) 99
8. J. A. Casas, F. Gomez and C. Munoz, CERN preprint, TH6194/91.
9. T. Kobayashi and N. Ohtsubu, Kanazawa preprint, DPKU–9103.
10. T. T. Burwick, R. K. Kaiser and H. F. Muller, Nucl. Phys. B355 (1991) 689
11. J. Erler, D. Jungnickel and J. Lauer, Phys. Rev D45 (1992) 3651; S. Stieberger, D. Jungnickel, J. Lauer and M. Spalinski, preprint, MPI–Ph/92–24; J. Erler, D. Jungnickel, M. Spalinski and S. Stieberger, preprint, MPI–Ph/92–56.
12. D. Bailin, A. Love and W. A. Sabra, Mod. Phys. Lett A6 (1992) 3607.
13. S. Stieberger, preprint, TUM–TH–151/92.
14. V. S. Kaplunovsky, Nucl. Phys. B307 (1988) 145; L. J. Dixon, V. S. Kaplunovsky and J. Louis, Nucl. Phys. B355 (1991) 649; J. P. Derendinger, S. Ferrara, C. Kounas and F. Zwirner, Nucl. Phys. B372 (1992) 145, Phys. Lett. B271 (1991) 307.

15. I. Antoniadis, J. Ellis, R. Lacaze and D. V. Nanopoulos, CERN–TH 6136/91; S. Kalara, J. L. Lopez and D. V. Nanopoulos, Phys. Lett. B269 (1991) 84.
16. L. E. Ibanez and D. Lüst and G. G. Ross, Phys. Lett. B272 (1991) 251; D. Bailin and A. Love, Phys. Lett. B278 (1992) 125.
17. L. E. Ibanez and D. Lüst, CERN–TH 6380/92.
18. D. Bailin and A. Love, Phys. Lett. B292 (1992) 315.
19. Y. Katsuki, Y. Kawanura, T. Kobayashi and N. Ohtsubo,Y. Ono and K. Tanioka, Nucl. Phys. B341 (1990) 611; D. Narkushevich, N. Olshanetsky and A. Perelonov, Comm. Math. Phys. 111 (1987) 247.
20. T. Kobayashi and N. Ohtsubo, Phys. Lett. B262 (1991) 425.
21. D. J. Gross, J. A. Harvey, E. Martinec and R. Rohm, Nucl. Phys. B267 (1986) 75.
22. A. Font, L. E. Ibanez and F. Quevedo, Phys. Lett. B217 (1989) 272.
23. T. Kobayashi and N. Ohtsubo, Phys. Lett. B245 (1990) 441.

Table Captions

Table 1: Point group elements and lattices for $\mathbf{Z}_M \times \mathbf{Z}_N$ orbifolds. The point group elements θ and ω which are of the form $(e\lambda 2\pi i\eta_1, e\lambda 2\pi i\eta_2, e\lambda 2\pi i\eta_3)$, are specified in the table by (η_1, η_2, η_3).

Table 2: Fixed points (and tori) for the $\mathbf{Z}_2 \times \mathbf{Z}_6$ orbifold and associated space group elements. The inequivalent fixed points (or tori) are obtained by giving integers n with half integral coefficients the values $0, 1$ and integers n with third integral coefficients the values $0, \pm 1$. The coefficients $a_1, ..., a_6$ are arbitrary, and for the space group elements (α, l) the lattice vector $(I - \alpha)\Lambda$, where Λ is an arbitrary lattice vector, may be added to l.

Table 3: Action of θ and ω on the fixed tori for the $\mathbf{Z}_2 \times \mathbf{Z}_6$ orbifold.

Table 4: Suppression factors $e\lambda{-}\lambda_1 R_1 \lambda 2/4\pi\sqrt{3} e\lambda{-}\lambda_3 R_3 \lambda 2/4\pi\sqrt{3} e\lambda{-}\lambda_5 R_5 \lambda 2/4\pi\sqrt{3}$ for the $\mathbf{Z}_2 \times \mathbf{Z}_6$ orbifold Yukawa couplings. The values of the integers characterizing fixed points (or tori) for which non-trivial suppression factors occur are indicated. The $\theta\lambda\alpha\omega\lambda\beta$ twisted sector is denoted by $T_{\alpha\beta}$.

TABLE 1

Point Group	θ	ω	Lattice
$\mathbf{Z}_2 \times \mathbf{Z}_2$	$(1,0,-1)/2$	$(0,1,-1)/2$	$SO(4)\lambda 3$
$\mathbf{Z}_3 \times \mathbf{Z}_3$	$(1,0,-1)/3$	$(0,1,-1)/3$	$SU(3)\lambda 3$
$\mathbf{Z}_2 \times \mathbf{Z}_4$	$(1,0,-1)/2$	$(0,1,-1)/4$	$SO(4) \times SO(5)\lambda 2$
$\mathbf{Z}_4 \times \mathbf{Z}_4$	$(1,0,-1)/4$	$(0,1,-1)/4$	$SO(5)\lambda 3$
$\mathbf{Z}_2 \times \mathbf{Z}_6$	$(1,0,-1)/2$	$(0,1,-1)/6$	$SO(4) \times G_2\lambda 2$
$\mathbf{Z}_2 \times \mathbf{Z}_6'$	$(1,0,-1)/2$	$(1,1,-2)/6$	$G_2\lambda 3$
$\mathbf{Z}_3 \times \mathbf{Z}_6$	$(1,0,-1)/3$	$(0,1,-1)/6$	$SU(3) \times G_2\lambda 2$
$\mathbf{Z}_6 \times \mathbf{Z}_6$	$(1,0,-1)/6$	$(0,1,-1)/6$	$G_2\lambda 3$

TABLE 2

twisted sector	Fixed point or torus	l
θ	$\frac{n_1}{2}e_1 + \frac{n_2}{2}e_2 + \frac{n_5}{2}e_5 + \frac{n_6}{2}e_6 + a_3e_3 + a_4e_4$	$n_1e_1 + n_2e_2 + n_5e_5 + n_6e_6$
ω	$a_1e_1 + a_2e_2$	0
$\omega\lambda 2$	$-\frac{n_3}{3}e_4 + \frac{n_5}{3}e_6 + a_1e_1 + a_2e_2$	$n_3e_3 + n_5e_5$
$\omega\lambda 3$	$\frac{n_3}{2}e_3 + \frac{n_4}{2}e_4 + \frac{n_5}{2}e_5 + \frac{n_6}{2}e_6 + a_1e_1 + a_2e_2$	$n_3e_3 + n_4e_4 + n_5e_5 + n_6e_6$
$\omega\lambda 4$	$-\frac{n_3}{3}e_4 + \frac{n_5}{3}e_6 + a_1e_1 + a_2e_2$	$-n_3e_3 - n_5e_5$
$\omega\lambda 5$	$a_1e_1 + a_2e_2$	0
$\theta\omega$	$-\frac{n_1}{2}e_1 - \frac{n_2}{2}e_2 - \frac{n_5}{3}e_6$	$n_1e_1 + n_2e_2 + n_5e_5$
$\theta\omega\lambda 2$	$-\frac{n_1}{2}e_1 - \frac{n_2}{2}e_2 - \frac{n_3}{3}e_4$	$n_1e_1 + n_2e_2 + n_3e_3$
$\theta\omega\lambda 3$	$\frac{n_1}{2}e_1 + \frac{n_2}{2}e_2 + \frac{n_3}{2}e_3 + \frac{n_4}{2}e_4 + a_5e_5 + a_6e_6$	$n_1e_1 + n_2e_2 + n_3e_3 + n_4e_4$

TABLE 3

twisted sector	Action of θ	Action of ω
θ	I	$(n_5, n_6) \rightarrow (n_6, n_5 + n_6)$
ω	I	I
$\omega\lambda 2$	$n_5 \rightarrow -n_5$	$(n_3, n_5) \rightarrow (-n_3, -n_5)$
$\omega\lambda 3$	I	$(n_3, n_4, n_5, n_6) \rightarrow (n_3 + n_4, n_3, n_6, n_5 + n_6)$
$\omega\lambda 4$	$n_5 \rightarrow -n_5$	$(n_3, n_5) \rightarrow (-n_3, -n_5)$
$\omega\lambda 5$	I	I
$\theta\omega$	$n_5 \rightarrow -n_5$	I
$\theta\omega\lambda 2$	I	$n_3 \rightarrow -n_3$
$\theta\omega\lambda 3$	I	$(n_3, n_4) \rightarrow (n_4, n_3 + n_4)$

TABLE 4

Yukawa Coupling	λ_1	λ_2	λ_3
$T_{01}T_{12}T_{13}$	0	$\mu(n_3, p_3, p_4)$	0
$T_{02}T_{11}T_{13}$	0	$\mu(n_3, p_3, p_4)$	0
$T_{03}T_{10}T_{13}$	0	0	0
$T_{04}T_{10}T_{12}$	0	0	$\mu(n_5, p_5, p_6)$
$T_{04}T_{11}T_{11}$	0	1 for $n_3(T_{04}) \neq 0$	2 for $m_5(T_{11}) - n_5(T_{11}) \neq 0$
$T_{05}T_{10}T_{11}$	0	0	$\mu(n_5, p_5, p_6)$

APPENDIX

$\mathbf{Z}_2 \times \mathbf{Z}_2$ *Orbifold*

Lattice : $SO(4)\lambda 3$

$$\theta = (C(SO(4)), I, C(SO(4))), \quad \omega = (I, C(SO(4)), C(SO(4))).$$

The fixed tori and associated space group elements are given with the conventions of table 2 by

twisted sector	Fixed point or torus	l
θ	$\frac{n_1}{2}e_1 + \frac{n_2}{2}e_2 + \frac{n_5}{2}e_5 + \frac{n_6}{2}e_6 + a_3e_3 + a_4e_4$	$n_1e_1 + n_2e_2 + n_5e_5 + n_6e_6$
ω	$\frac{n_3}{2}e_3 + \frac{n_4}{2}e_4 + \frac{n_5}{2}e_5 + \frac{n_6}{2}e_6 + a_1e_1 + a_2e_2$	$n_3e_3 + n_4e_4 + n_5e_5 + n_6e_6$
$\theta\omega$	$\frac{n_1}{2}e_1 + \frac{n_2}{2}e_2 + \frac{n_3}{2}e_3 + \frac{n_4}{2}e_4 + a_5e_5 + a_6e_6$	$n_1e_1 + n_2e_2 + n_3e_3 + n_4e_4$

The only non-zero Yukawa couplings is $T_{01}T_{10}T_{11}$, where $T_{\alpha\beta}$ denotes the $\theta\lambda\alpha\omega\lambda\beta$ twisted sector, and since no complex plane is twisted by all three twist fields there is no moduli dependence.

$\mathbf{Z}_3 \times \mathbf{Z}_3$ *Orbifold*

Lattice $SU(3)\lambda 3$

Already discussed in [12].

$\mathbf{Z}_2 \times \mathbf{Z}_4$ *Orbifold*

lattice $SO(4) \times SO(5) \times SO(5)$

In the orthonormal basis,

$$\begin{aligned} \theta X_1 &= -X_1, & \omega X_1 &= X_1, \\ \theta X_2 &= X_2, & \omega X_2 &= e\lambda 2\pi i/4 X_2, \\ \theta X_3 &= -X_3, & \omega X_3 &= e\lambda{-2}\pi i/4 X_3. \end{aligned}$$

In the lattice basis,

$$\theta = (C(SO(4)), I, C\lambda 2(SO(5))), \quad \omega = (I, C(SO(5)), C\lambda{-1}(SO(5)))$$

so that

$$\begin{aligned} \theta e_1 &= -e_1, & \theta e_3 &= e_3, & \theta e_5 &= -e_5, \\ \theta e_2 &= -e_2, & \theta e_4 &= e_4, & \theta e_6 &= -e_6 \end{aligned}$$

and

$$\begin{aligned} \omega e_1 &= e_1, & \omega e_3 &= e_3 + 2e_4, & \omega e_5 &= -e_5 - 2e_6, \\ \omega e_2 &= e_2, & \omega e_4 &= -e_3 - e_4, & \omega e_6 &= e_5 + e_6. \end{aligned}$$

The independent deformation parameters (moduli) are R_1, R_2, $\cos\theta_{12}$, R_3 and R_5, where

$$|e_1| \equiv R_1, \quad |e_2| \equiv R_2, \quad |e_3| = \sqrt{2}|e_4| \equiv R_3, \quad |e_5| = \sqrt{2}|e_6| \equiv R_5,$$

also,

$$\cos\theta_{34} = \cos\theta_{56} = -\frac{1}{\sqrt{2}}$$

with all other $\cos\theta_{ij}$ except $\cos\theta_{12}$ zero. A suitable choice of orthonormal basis $\tilde{e}_1, ..., \tilde{e}_6$ is given by

$$\begin{aligned} e_1 &= R_1\tilde{e}_1, & e_3 &= R_3\tilde{e}_3, & e_5 &= R_5\tilde{e}_5, \\ e_2 &= R_2(\cos\theta_{12}\tilde{e}_1 + \sin\theta_{12}\tilde{e}_2), & e_4 &= \frac{R_3}{2}(-\tilde{e}_3 + \tilde{e}_4), & e_6 &= \frac{R_5}{2}(-\tilde{e}_5 + \tilde{e}_6). \end{aligned}$$

The fixed points or tori and associated space group elements are given with the conventions of table 2 by

twisted sector	Fixed point or torus	l
θ	$\frac{n_1}{2}e_1+\frac{n_2}{2}e_2+\frac{n_5}{2}e_5+\frac{n_6}{2}e_6+a_3e_3+a_4e_4$	$n_1e_1+n_2e_2+n_5e_5+n_6e_6$
ω	$a_1e_1+a_2e_2-\frac{n_4}{2}e_3+\frac{n_6}{2}e_5$	$n_4e_4-n_6e_6$
$\omega\lambda 2$	$a_1e_1+a_2e_2+\frac{n_3}{2}e_3+\frac{n_4}{2}e_4+\frac{n_5}{2}e_5+\frac{n_6}{2}e_6$	$n_3e_3+n_4e_4+n_5e_5+n_6e_6$
$\omega\lambda 3$	$a_1e_1+a_2e_2-\frac{n_4}{2}e_3+\frac{n_6}{2}e_5$	$-n_4e_4+n_6e_6$
$\theta\omega$	$\frac{n_1}{2}e_1+\frac{n_2}{2}e_2-\frac{n_4}{2}e_3+\frac{n_6}{2}e_5$	$n_1e_1+n_2e_2+n_4e_4+n_6e_6$
$\theta\omega\lambda 2$	$\frac{n_1}{2}e_1+\frac{n_2}{2}e_2+\frac{n_3}{2}e_3+\frac{n_4}{2}e_4+a_5e_5+a_6e_6$	$n_1e_1+n_2e_2+n_3e_3+n_4e_4$
$\theta\omega\lambda 3$	$\frac{n_1}{2}e_1+\frac{n_2}{2}e_2-\frac{n_4}{2}e_3+\frac{n_6}{2}e_5$	$n_1e_1+n_2e_2+n_6e_6-n_4e_4$

The action of θ and ω on the fixed points (or tori) is given by

Twisted sector	Action of θ	Action of ω
θ	I	$n_5 \to n_5+n_6$
$\omega, \omega\lambda 3$	I	I
$\omega\lambda 2$	I	$(n_3,n_5) \to (n_3+n_4, n_5+n_6)$
$\theta\omega, \theta\omega\lambda 3$	I	I
$\theta\omega\lambda 2$	I	$n_3 \to n_3+n_4$

The moduli dependent suppression factors for the non-zero Yukawa couplings amongst twisted sectors containing massless states written in the form $e\lambda-\lambda_{2i-1}R_{2i-1}\lambda 2/16\pi$█ are given by

Yukawa Coupling	λ_1	λ_2	λ_3
$T_{01}T_{11}T_{12}$	0	$\rho(n_4,p_3,p_4)$	0
$T_{02}T_{10}T_{12}$	0	0	0
$T_{03}T_{10}T_{11}$	0	0	$\rho(n_6,p_5,p_6)$
$T_{11}T_{11}T_{02}$	0	$\rho(n_4,p_3,p_4)$	$\rho(n_6,p_5,p_6)$

$\mathbf{Z}_4 \times \mathbf{Z}_4$ *Orbifold*

Lattice $SO(5) \times SO(5) \times SO(5)$

In the orthonormal basis,

$$\begin{aligned}
\theta X_1 &= e\lambda 2\pi i/4 X_1, & \omega X_1 &= X_1, \\
\theta X_2 &= X_2, & \omega X_2 &= e\lambda 2\pi i/4 X_2, \\
\theta X_3 &= e\lambda{-}2\pi i/4 X_3, & \omega X_3 &= e\lambda{-}2\pi i/4 X_3.
\end{aligned}$$

In the lattice basis,

$$\theta = (C(SO(5)), I, C\lambda{-}1(SO(5))), \quad \omega = (I, C(SO(5)), C\lambda{-}1(SO(5)))$$

so that

$$\begin{aligned}
\theta e_1 &= e_1 + 2e_2, & \theta e_3 &= e_3, & \theta e_5 &= -e_5 - 2e_6, \\
\theta e_2 &= -e_1 - e_2, & \theta e_4 &= e_4, & \theta e_6 &= e_5 + e_6
\end{aligned}$$

and

$$\begin{aligned}
\omega e_1 &= e_1, & \omega e_3 &= e_3 + 2e_4, & \omega e_5 &= -e_5 - 2e_6, \\
\omega e_2 &= e_2, & \omega e_4 &= -e_3 - e_4, & \omega e_6 &= e_5 + e_6.
\end{aligned}$$

The independent deformation parameters (moduli) are R_1, R_3 and R_5, where

$$|e_1| = \sqrt{2}|e_2| \equiv R_1, \quad |e_3| = \sqrt{2}|e_4| \equiv R_3, \quad |e_5| = \sqrt{2}|e_6| \equiv R_5,$$

also

$$\cos\theta_{12} = \cos\theta_{34} = \cos\theta_{56} = -\frac{1}{\sqrt{2}}$$

with all other $\cos\theta_{ij}$ zero. A suitable choice of orthonormal basis $\tilde{e}_1, ..., \tilde{e}_6$ is given by

$$\begin{aligned}
e_1 &= R_1\tilde{e}_1, & e_3 &= R_3\tilde{e}_3, & e_5 &= R_5\tilde{e}_5, \\
e_2 &= -\frac{R_1}{2}(\tilde{e}_1 - \tilde{e}_2), & e_4 &= -\frac{R_3}{2}(\tilde{e}_3 - \tilde{e}_4), & e_6 &= -\frac{R_5}{2}(\tilde{e}_5 - \tilde{e}_6).
\end{aligned}$$

The fixed points or tori and associated space group elements are given with the conventions of table 2 by

twisted sector	Fixed point or torus	l
θ	$-\frac{n_2}{2}e_1+\frac{n_6}{2}e_5+a_3e_3+a_4e_4$	$n_2e_2+n_6e_6$
ω	$a_1e_1+a_2e_2-\frac{n_4}{2}e_3+\frac{n_6}{2}e_5$	$n_4e_4+n_6e_6$
$\omega\lambda2$	$a_1e_1+a_2e_2+\frac{n_3}{2}e_3+\frac{n_4}{2}e_4+\frac{n_5}{2}e_5+\frac{n_6}{2}e_6$	$n_3e_3+n_4e_4+n_5e_5+n_6e_6$
$\omega\lambda3$	$a_1e_1+a_2e_2+\frac{n_4}{2}e_3-\frac{n_6}{2}e_5$	$n_4e_4+n_6e_6$
$\theta\omega$	$-\frac{n_2}{2}e_1-\frac{n_4}{2}e_3+\frac{n_5}{2}e_5+\frac{n_6}{2}e_6$	$n_2e_2+n_4e_4+n_5e_5+n_6e_6$
$\theta\omega\lambda2$	$-\frac{n_2}{2}e_1+\frac{n_3}{2}e_3+\frac{n_4}{2}e_4-\frac{n_6}{2}e_5$	$n_2e_2+n_4e_4+n_3e_3+n_6e_6$
$\theta\omega\lambda3$	$-\frac{n_2}{2}e_1+\frac{n_4}{2}e_3+a_5e_5+a_6e_6$	$n_2e_2+n_4e_4$
$\theta\lambda2$	$\frac{n_1}{2}e_1+\frac{n_2}{2}e_2+a_3e_3+a_4e_4+\frac{n_5}{2}e_5+\frac{n_6}{2}e_6$	$n_1e_1+n_2e_2+n_5e_5+n_6e_6$
$\theta\lambda3$	$\frac{n_2}{2}e_1+a_3e_3+a_4e_4-\frac{n_6}{2}e_5$	$n_2e_2+n_6e_6$
$\theta\lambda2\omega$	$\frac{n_1}{2}e_1+\frac{n_2}{2}e_2-\frac{n_4}{2}e_3-\frac{n_6}{2}e_5$	$n_1e_1+n_2e_2+n_4e_4+n_6e_6$
$\theta\lambda2\omega\lambda2$	$\frac{n_1}{2}e_1+\frac{n_2}{2}e_2+\frac{n_3}{2}e_3+\frac{n_4}{2}e_4+a_5e_5+a_6e_6$	$n_1e_1+n_2e_2+n_3e_3+n_4e_4$
$\theta\lambda3\omega$	$\frac{n_2}{2}e_1-\frac{n_4}{2}e_3+a_5e_5+a_6e_6$	$n_2e_2+n_4e_4$
$\theta\lambda2\omega\lambda3$	$\frac{n_1}{2}e_1+\frac{n_2}{2}e_2+\frac{n_4}{2}e_3+\frac{n_6}{2}e_5$	$n_1e_1+n_2e_2+n_4e_4+n_6e_6$
$\theta\lambda3\omega\lambda2$	$\frac{n_2}{2}e_1+\frac{n_3}{2}e_3+\frac{n_4}{2}e_4+\frac{n_6}{2}e_5$	$n_2e_2+n_3e_3+n_4e_4+n_6e_6$
$\theta\lambda3\omega\lambda3$	$\frac{n_2}{2}e_1+\frac{n_4}{2}e_3+\frac{n_5}{2}e_5+\frac{n_6}{2}e_6$	$n_2e_2+n_4e_4+n_5e_5+n_6e_6$

The action of θ and ω on the fixed points (or tori) is given by

twisted sector	Action of theta	Action of omega
$\theta, \theta\lambda 3$	I	I
$\omega, \omega\lambda 3$	I	I
$\theta\lambda 3\omega, \theta\omega\lambda 3$	I	I
$\theta\omega, \theta\lambda 3\omega\lambda 3$	$n_5 \rightarrow n_5 + n_6$	$n_5 \rightarrow n_5 + n_6$
$\theta\omega\lambda 2, \theta\lambda 3\omega\lambda 2$	I	$n_3 \rightarrow n_3 + n_4$
$\theta\lambda 2\omega, \theta\lambda 2\omega\lambda 3$	$n_1 \rightarrow n_1 + n_2$	I
$\omega\lambda 2$	$n_5 \rightarrow n_5 + n_6$	$(n_3, n_5) \rightarrow (n_3 + n_4, n_5 + n_6)$
$\theta\lambda 2$	$(n_1, n_5) \rightarrow (n_1 + n_2, n_5 + n_6)$	$n_5 \rightarrow n_5 + n_6$
$\theta\lambda 2\omega\lambda 2$	$n_1 \rightarrow n_1 + n_2$	$n_3 \rightarrow n_3 + n_4$

The moduli dependent suppression factors for the non-zero Yukawa couplings amongst twisted sectors containing massless states written in the form $e\lambda-\lambda_{2i-1}R_{2i-1}\lambda 2/16\pi$ are given by

Yukawa coupling	λ_1	λ_2	λ_3
$T_{01}T_{12}T_{31}$	0	$\rho(n_4, p_3, p_4)$	0
$T_{01}T_{13}T_{30}$	0	0	0
$T_{01}T_{21}T_{22}$	0	$\rho(n_4, p_3, p_4)$	0
$T_{02}T_{11}T_{31}$	0	$\rho(n_4, p_3, p_4)$	0
$T_{02}T_{12}T_{30}$	0	0	$\rho(n_6, p_5, p_6)$
$T_{02}T_{20}T_{22}$	0	0	0
$T_{02}T_{21}T_{21}$	0	$\rho(n_4, p_3, p_4)$	$\rho(n_6, p_5, p_6)$
$T_{03}T_{10}T_{31}$	0	0	0
$T_{03}T_{11}T_{30}$	0	0	$\rho(n_6, p_5, p_6)$
$T_{03}T_{20}T_{21}$	0	0	$\rho(n_6, p_5, p_6)$
$T_{10}T_{12}T_{22}$	$\rho(n_2, p_1, p_2)$	0	0
$T_{10}T_{13}T_{21}$	$\rho(n_2, p_1, p_2)$	0	0
$T_{11}T_{12}T_{21}$	$\rho(n_2, p_1, p_2)$	$\rho(n_4, p_3, p_4)$	$\rho(n_6, p_5, p_6)$
$T_{11}T_{11}T_{22}$	$\rho(n_2, p_1, p_2)$	$\rho(n_4, p_3, p_4)$	0
$T_{11}T_{13}T_{20}$	$\rho(n_2, p_1, p_2)$	0	0
$T_{12}T_{12}T_{20}$	$\rho(n_2, p_1, p_2)$	0	$\rho(n_6, p_5, p_6)$

$\mathbf{Z}_2 \times \mathbf{Z}_6$ *Orbifold*

Lattice $SO(4) \times G_2\lambda 2$

Discussed in section 3.

$\mathbf{Z}_2 \times \mathbf{Z}_6'$ *Orbifold*

Lattice $G_2\lambda 3$

In the orthonormal basis,

$$\begin{aligned} \theta X_1 &= -X_1, & \omega X_1 &= e\lambda 2\pi i/6 X_1, \\ \theta X_2 &= X_2, & \omega X_2 &= e\lambda 2\pi i/6 X_2, \\ \theta X_3 &= -X_3, & \omega X_3 &= e\lambda{-2}\pi i/3 X_3. \end{aligned}$$

In the lattice basis,

$$\theta = (C\lambda 3(G_2), I, C\lambda 3(G_2)), \quad \omega = (C(G_2), C(G_2), C\lambda{-2}(G_2))$$

so that

$$\begin{aligned} \theta e_1 &= -e_1, & \theta e_3 &= e_3, & \theta e_5 &= -e_5, \\ \theta e_2 &= -e_2, & \theta e_4 &= e_4, & \theta e_6 &= -e_6. \end{aligned}$$

and

$$\begin{aligned} \omega e_1 &= -e_1 - e_2, & \omega e_3 &= -e_3 - e_4, & \omega e_5 &= e_5 + e_6, \\ \omega e_2 &= 3e_1 + 2e_2, & \omega e_4 &= 3e_3 + 2e_4, & \omega e_6 &= -3e_5 - 2e_6. \end{aligned}$$

The independent deformation parameters (moduli) are R_1, R_3 and R_5, where

$$|e_2| = \sqrt{3}|e_1| \equiv \sqrt{3}R_1, \quad |e_4| = \sqrt{3}|e_3| \equiv \sqrt{3}R_3, \quad |e_6| = \sqrt{3}|e_5| \equiv \sqrt{3}R_5$$

also,

$$\cos\theta_{12} = \cos\theta_{34} = \cos\theta_{56} = -\frac{\sqrt{3}}{2}$$

with all other $\cos\theta_{ij}$ zero. A suitable choice of orthonormal basis $\tilde{e}_1, ..., \tilde{e}_6$ is given by

$$e_1 = R_1\tilde{e}_1 \qquad e_3 = R_3\tilde{e}_3 \qquad e_5 = R_5\tilde{e}_5$$

$$e_2 = R_1(-\frac{3}{2}\tilde{e}_1 - \frac{\sqrt{3}}{2}\tilde{e}_2) \qquad e_4 = R_3(-\frac{3}{2}\tilde{e}_3 - \frac{\sqrt{3}}{2}\tilde{e}_4) \qquad e_6 = R_5(-\frac{3}{2}\tilde{e}_5 - \frac{\sqrt{3}}{2}\tilde{e}_6)$$

The fixed points or tori and associated space group elements are given with the conventions of table 2 by

Twisted sector	Fixed point or torus	l
θ	$\frac{n_1}{2}e_1 + \frac{n_2}{2}e_2 + \frac{n_5}{2}e_5 + \frac{n_6}{2}e_6 + a_3e_3 + a_4e_4$	$n_1e_1 + n_2e_2 + n_5e_5 + n_6e_6$
ω	$\frac{n_5}{3}e_6$	n_5e_5
$\omega\lambda 2$	$-\frac{n_1}{3}e_2 - \frac{n_3}{3}e_4 + \frac{n_5}{3}e_6$	$n_1e_1 + n_3e_3 - n_5e_5$
$\omega\lambda 3$	$\frac{n_1}{2}e_1 + \frac{n_2}{2}e_2 + \frac{n_3}{2}e_3 + \frac{n_4}{2}e_4 + a_5e_5 + a_6e_6$	$n_1e_1 + n_2e_2 + n_3e_3 + n_4e_4$
$\omega\lambda 4$	$-\frac{n_1}{3}e_2 - \frac{n_3}{3}e_4 + \frac{n_5}{3}e_6$	$-n_1e_1 - n_3e_3 + n_5e_5$
$\omega\lambda 5$	$\frac{n_5}{3}e_6$	$-n_5e_5$
$\theta\omega$	$\frac{n_1}{3}e_1$	n_1e_1
$\theta\omega\lambda 2$	$-\frac{n_3}{3}e_4$	n_3e_3
$\theta\omega\lambda 3$	$\frac{n_3}{2}e_3 + \frac{n_4}{2}e_4 + \frac{n_5}{2}e_5 + \frac{n_6}{2}e_6 + a_1e_1 + a_2e_2$	$n_3e_3 + n_4e_4 + n_5e_5 + n_6e_6$
$\theta\omega\lambda 4$	$-\frac{n_3}{3}e_4$	$-n_3e_3$
$\theta\omega\lambda 5$	$\frac{n_1}{3}e_2$	$-n_1e_1$

The action of θ and ω on the fixed points (or tori) is given by

Twisted sector	Action of θ	Action of ω
θ	I	$(n_1, n_2, n_5, n_6) \to (n_1 + n_2, n_1, n_5 + n_6, n_5)$
$\omega, \omega\lambda5$	$n_5 \to -n_5$	I
$\omega\lambda2, \omega\lambda4$	$(n_3, n_5) \to (-n_3, -n_5)$	$(n_1, n_3) \to (-n_1, -n_3)$
$\omega\lambda3$	I	$(n_1, n_2, n_3, n_4) \to (n_1 + n_2, n_1, n_3 + n_4, n_3)$
$\theta\omega, \theta\omega\lambda5$	$n_1 \to -n_1$	$n_1 \to -n_1$
$\theta\omega\lambda2, \theta\omega\lambda4$	I	$n_3 \to -n_3$
$\theta\omega\lambda3$	I	$(n_3, n_4, n_5, n_6) \to (n_3 + n_4, n_3, n_5 + n_6, n_5)$

The moduli dependent suppression factors for the non-zero Yukawa couplings amongst twisted sectors containing massless states are given in the conventions of table 4 by

Yukawa Coupling	λ_1	λ_2	λ_3
$T_{01}T_{02}T_{03}$	$\mu(n_1, p_1, p_2)$	$\mu(n_3, p_3, p_4)$	0
$T_{01}T_{11}T_{14}$	1 for $n_1(T_{11}) \neq 0$	1 for $n_3(T_{14}) \neq 0$	1 for $n_5(T_{01}) \neq 0$
$T_{02}T_{02}T_{02}$	2 for $m_1(T_{02}) - n_1(T_{02}) \neq 0$	2 for $m_3(T_{02}) - n_3(T_{02}) \neq 0$	2 for $m_5(T_{02}) - n_5(T_{02}) \neq 0$
$T_{02}T_{10}T_{14}$	$\mu(n_1, p_1, p_2)$	0	$\mu(n_5, p_5, p_6)$
$T_{02}T_{11}T_{13}$	0	$\mu(n_3, p_3, p_4)$	$\mu(n_5, p_5, p_6)$
$T_{03}T_{10}T_{13}$	0	0	0

$\mathbf{Z}_3 \times \mathbf{Z}_6$ *Orbifold*

lattice $SU(3) \times G_2\lambda 2$

In the orthonormal basis,

$$\begin{aligned} \theta X_1 &= e\lambda 2\pi i/3X_1, & \omega X_1 &= X_1, \\ \theta X_2 &= X_2, & \omega X_2 &= e\lambda 2\pi i/6X_2, \\ \theta X_3 &= e\lambda{-}2\pi i/3X_3, & \omega X_3 &= e\lambda{-}2\pi i/6X_3. \end{aligned}$$

In the lattice basis,

$$\theta = (C(SU(3)), I, C\lambda{-}2(G_2)), \quad \omega = (I, C(G_2), C\lambda{-}1(G_2))$$

so that

$$\begin{aligned} \theta e_1 &= e_2, & \theta e_3 &= e_3, & \theta e_5 &= e_5 + e_6, \\ \theta e_2 &= -e_1 - e_2, & \theta e_4 &= e_4, & \theta e_6 &= -3e_5 - 2e_6. \end{aligned}$$

and

$$\begin{aligned} \omega e_1 &= e_1, & \omega e_3 &= -e_3 - e_4, & \omega e_5 &= 2e_5 + e_6, \\ \omega e_2 &= e_2, & \omega e_4 &= 3e_3 + 2e_4, & \omega e_6 &= -3e_5 - e_6. \end{aligned}$$

The independent deformation parameters (moduli) are R_1, R_3 and R_5, where

$$|e_2| = |e_1| \equiv R_1, \quad |e_4| = \sqrt{3}|e_3| \equiv \sqrt{3}R_3, \quad |e_6| = \sqrt{3}|e_5| \equiv \sqrt{3}R_5$$

also,

$$\cos\theta_{12} = -\frac{1}{2}, \quad \cos\theta_{34} = \cos\theta_{56} = -\frac{\sqrt{3}}{2}$$

with all other $\cos\theta_{ij}$ zero. A suitable choice of orthonormal basis $\tilde{e}_1, ..., \tilde{e}_6$ is given by

$$\begin{aligned} e_1 &= R_1\tilde{e}_1, & e_3 &= R_3\tilde{e}_3, & e_5 &= R_5\tilde{e}_5, \\ e_2 &= R_1(-\frac{1}{2}\tilde{e}_1 + \frac{\sqrt{3}}{2}\tilde{e}_2), & e_4 &= R_3(-\frac{3}{2}\tilde{e}_3 - \frac{\sqrt{3}}{2}\tilde{e}_4), & e_6 &= R_5(-\frac{3}{2}\tilde{e}_5 - \frac{\sqrt{3}}{2}\tilde{e}_6). \end{aligned}$$

The fixed points or tori and associated group elements are given with the conventions of table 2 by

Twisted sector	Fixed point or torus	l
θ	$\frac{n_1}{3}(2e_1+e_2)+\frac{n_5}{3}e_6+a_3e_3+a_4e_4$	$n_1e_1+n_5e_5$
$\theta\lambda 2$	$\frac{n_1}{3}(2e_1+e_2)+\frac{n_5}{3}e_6+a_3e_3+a_4e_4$	$-n_1e_1-n_5e_5$
ω	$a_1e_1+a_2e_2$	0
$\omega\lambda 2$	$-\frac{n_3}{3}e_4+\frac{n_5}{3}e_6+a_1e_1+a_2e_2$	$n_3e_3+n_5e_5$
$\omega\lambda 3$	$\frac{n_3}{2}e_3+\frac{n_4}{2}e_4+\frac{n_5}{2}e_5+\frac{n_6}{2}e_6+a_1e_1+a_2e_2$	$n_3e_3+n_4e_4+n_5e_5+n_6e_6$
$\omega\lambda 4$	$-\frac{n_3}{3}e_4+\frac{n_5}{3}e_6+a_1e_1+a_2e_2$	$-n_3e_3-n_5e_5$
$\omega\lambda 5$	$a_1e_1+a_2e_2$	0
$\theta\omega$	$\frac{n_1}{3}(2e_1+e_2)+\frac{n_5}{2}e_5+\frac{n_6}{2}e_6$	$n_1e_1+n_5e_5+n_6e_6$
$\theta\omega\lambda 2$	$\frac{n_1}{3}(2e_1+e_2)-\frac{n_3}{3}e_4+\frac{n_5}{3}e_6$	$n_1e_1+n_3e_3-n_5e_5$
$\theta\omega\lambda 3$	$\frac{n_1}{3}(2e_1+e_2)+\frac{n_3}{2}e_3+\frac{n_4}{2}e_4$	$n_1e_1+n_3e_3+n_4e_4$
$\theta\omega\lambda 4$	$\frac{n_1}{3}(2e_1+e_2)-\frac{n_3}{2}e_4+a_5e_5+a_6e_6$	$n_1e_1-n_3e_3$
$\theta\omega\lambda 5$	$\frac{n_1}{3}(2e_1+e_2)$	n_1e_1
$\theta\lambda 2\omega$	$\frac{n_1}{3}(2e_1+e_2)$	$-n_1e_1$
$\theta\lambda 2\omega\lambda 2$	$\frac{n_1}{3}(2e_1+e_2)-\frac{n_3}{3}e_4$	$-n_1e_1+n_3e_3$
$\theta\lambda 2\omega\lambda 3$	$\frac{n_1}{3}(2e_1+e_2)+\frac{n_3}{2}e_3+\frac{n_4}{2}e_4$	$-n_1e_1+n_3e_3+n_4e_4$
$\theta\lambda 2\omega\lambda 4$	$\frac{n_1}{3}(2e_1+e_2)-\frac{n_3}{3}e_4+\frac{n_5}{3}e_6$	$-n_1e_1-n_3e_3+n_5e_5$
$\theta\lambda 2\omega\lambda 5$	$\frac{n_1}{3}(2e_1+e_2)+\frac{n_5}{2}e_5+\frac{n_6}{2}e_6$	$-n_1e_1+n_5e_5+n_6e_6$

The action of θ and ω on the fixed points (or tori) is given by

Twisted sector	Action of θ	Action of ω
$\theta, \theta\lambda 2$	I	$n_5 \to -n_5$
$\omega, \omega\lambda 5$	I	I
$\omega\lambda 2, \omega\lambda 4$	I	$(n_3, n_5) \to (-n_3, -n_5)$
$\omega\lambda 3$	$(n_5, n_6) \to (n_5 + n_6, n_5)$	$(n_3, n_4, n_5, n_6) \to (n_3 + n_4, n_3, n_6, n_5 + n_6)$
$\theta\omega, \theta\lambda 2\omega\lambda 5$	$(n_5, n_6) \to (n_5 + n_6, n_5)$	$(n_5, n_6) \to (n_6, n_5 + n_6)$
$\theta\omega\lambda 2, \theta\lambda 2\omega\lambda 4$	I	$(n_3, n_5) \to (-n_3, -n_5)$
$\theta\omega\lambda 3, \theta\lambda 2\omega\lambda 3$	I	$(n_3, n_4) \to (n_3 + n_4, n_3)$
$\theta\omega\lambda 4, \theta\lambda 2\omega\lambda 2$	I	$n_3 \to -n_3$
$\theta\omega\lambda 5, \theta\lambda 2\omega$	I	I

The moduli dependent suppression factors for the non-zero Yukawa couplings amongst twisted sectors containing massless states are given in the conventions of table 4 by

Yukawa Coupling	λ_1	λ_2	λ_3
$T_{01}T_{13}T_{22}$	0	$\mu(n_3, p_3, p_4)$	0
$T_{01}T_{14}T_{21}$	0	1 for $n_3(T_{14}) \neq 0$	0
$T_{02}T_{12}T_{22}$	0	2 for $m_3(T_{02}) - n_3(T_{12}) \neq 0$	0
$T_{02}T_{13}T_{21}$	0	$\mu(n_3, p_3, p_4)$	1 for $n_5(T_{02}) \neq 0$
$T_{02}T_{14}T_{20}$	0	0	0
$T_{03}T_{11}T_{22}$	0	$\mu(n_3, p_3, p_4)$	0
$T_{03}T_{12}T_{21}$	0	$\mu(n_3, p_3, p_4)$	$\mu(n_5, p_5, p_6)$
$T_{03}T_{13}T_{20}$	0	0	$\mu(n_5, p_5, p_6)$
$T_{04}T_{10}T_{22}$	0	0	0
$T_{04}T_{11}T_{21}$	0	1 for $n_3(T_{04}) \neq 0$	$\mu(n_5, p_5, p_6)$
$T_{04}T_{12}T_{20}$	0	0	2 for $m_5(T_{04}) - n_5(T_{12}) \neq 0$
$T_{05}T_{10}T_{21}$	0	0	1 for $n_5(T_{10}) \neq 0$
$T_{05}T_{11}T_{20}$	0	0	$\mu(n_5, p_5, p_6)$
$T_{10}T_{12}T_{14}$	2 for $m_1(T_{10}) - n_1(T_{12}) \neq 0$	0	0
$T_{10}T_{13}T_{13}$	2 for $m_1(T_{10}) - n_1(T_{13}) \neq 0$	0	1 for $n_5(T_{10}) \neq 0$
$T_{11}T_{11}T_{14}$	2 for $m_1(T_{11}) - n_1(T_{11}) \neq 0$	1 for $n_3(T_{14}) \neq 0$	0
$T_{11}T_{12}T_{13}$	2 for $m_1(T_{11}) - n_1(T_{12}) \neq 0$	$\mu(n_3, p_3, p_4)$	$\mu(n_5, p_5, p_6)$
$T_{12}T_{12}T_{12}$	2 for $m_1(T_{12}) - n_1(T_{12}) \neq 0$	2 for $m_3(T_{12}) - n_3(T_{12}) \neq 0$	2 for $m_5(T_{12}) - n_5(T_{12}) \neq 0$

$\mathbf{Z}_6 \times \mathbf{Z}_6$ *Orbifold*

Lattice $G_2\lambda 3$

In the orthonormal basis,

$$\begin{aligned} \theta X_1 &= e\lambda 2\pi i/6 X_1, & \omega X_1 &= X_1, \\ \theta X_2 &= X_2, & \omega X_2 &= e\lambda 2\pi i/6 X_2, \\ \theta X_3 &= e\lambda{-}2\pi i/6 X_3, & \omega X_3 &= e\lambda{-}2\pi i/6 X_3. \end{aligned}$$

In the lattice basis,

$$\theta = (C(G_2), I, C\lambda{-}1(G_2)), \quad \omega = (I, C(G_2), C\lambda{-}1(G_2))$$

so that

$$\begin{aligned} \theta e_1 &= -e_1 - e_2, & \theta e_3 &= e_3, & \theta e_5 &= 2e_5 + e_6, \\ \theta e_2 &= 3e_1 + 2e_2, & \theta e_4 &= e_4 & \theta e_6 &= -3e_5 - e_6 \end{aligned}$$

and

$$\begin{aligned} \omega e_1 &= e_1, & \omega e_3 &= -e_3 - e_4, & \omega e_5 &= 2e_5 + e_6, \\ \omega e_2 &= e_2, & \omega e_4 &= 3e_3 + 2e_4, & \omega e_6 &= -3e_5 - e_6. \end{aligned}$$

The independent deformation parameters (moduli) are R_1, R_3 and R_5, where

$$|e_2| = \sqrt{3}|e_1| \equiv \sqrt{3}R_1, \quad |e_4| = \sqrt{3}|e_3| \equiv \sqrt{3}R_3, \quad |e_6| = \sqrt{3}|e_5| \equiv \sqrt{3}R_5$$

also,

$$\cos\theta_{12} = \cos\theta_{34} = \cos\theta_{56} = -\frac{\sqrt{3}}{2}$$

with all other $\cos\theta_{ij}$ zero. A suitable choice of orthonormal basis $\tilde{e}_1, ..., \tilde{e}_6$ is given by

$$\begin{aligned} e_1 &= R_1\tilde{e}_1, & e_3 &= R_3\tilde{e}_3, & e_5 &= R_5\tilde{e}_5, \\ e_2 &= R_1(-\frac{3}{2}\tilde{e}_1 - \frac{\sqrt{3}}{2}\tilde{e}_2), & e_4 &= R_3(-\frac{3}{2}\tilde{e}_3 - \frac{\sqrt{3}}{2}\tilde{e}_4), & e_6 &= R_5(-\frac{3}{2}\tilde{e}_5 - \frac{\sqrt{3}}{2}\tilde{e}_6). \end{aligned}$$

The fixed points or tori and associated space group elements are given with the conventions of table 2 by

Twisted sector	Fixed point or torus	l
θ	$a_3e_3+a_4e_4$	0
$\theta\lambda 2$	$-\frac{n_1}{3}e_2+\frac{n_5}{3}e_6+a_3e_3+a_4e_4$	$n_1e_1+n_5e_5$
$\theta\lambda 3$	$\frac{n_1}{2}e_1+\frac{n_2}{2}e_2+\frac{n_5}{2}e_5+\frac{n_6}{2}e_6+a_3e_3+a_4e_4$	$n_1e_1+n_2e_2+n_5e_5+n_6e_6$
$\theta\lambda 4$	$-\frac{n_1}{3}e_2+\frac{n_5}{3}e_6+a_3e_3+a_4e_4$	$-n_1e_1-n_5e_5$
$\theta\lambda 5$	$a_3e_3+a_4e_4$	0
ω	$a_1e_1+a_2e_2$	0
$\omega\lambda 2$	$-\frac{n_3}{3}e_4+\frac{n_5}{3}e_6+a_1e_1+a_2e_2$	$n_3e_3+n_5e_5$
$\omega\lambda 3$	$\frac{n_3}{2}e_3+\frac{n_4}{2}e_4+\frac{n_5}{2}e_5+\frac{n_6}{2}e_6+a_1e_1+a_2e_2$	$n_3e_3+n_4e_4+n_5e_5+n_6e_6$
$\omega\lambda 4$	$-\frac{n_3}{3}e_4+\frac{n_5}{3}e_6+a_1e_1+a_2e_2$	$-n_3e_3-n_5e_5$
$\omega\lambda 5$	$a_1e_1+a_2e_2$	0
$\theta\omega$	$\frac{n_5}{3}e_6$	n_5e_5
$\theta\omega\lambda 2$	$-\frac{n_3}{3}e_4+\frac{n_5}{2}e_5+\frac{n_6}{2}e_6$	$n_3e_3+n_5e_5+n_6e_6$
$\theta\omega\lambda 3$	$\frac{n_3}{2}e_3+\frac{n_4}{2}e_4+\frac{n_5}{3}e_6$	$n_3e_3+n_4e_4-n_5e_5$
$\theta\omega\lambda 4$	$-\frac{n_3}{3}e_4$	$-n_3e_3$
$\theta\omega\lambda 5$	$a_5e_5+a_6e_6$	0
$\theta\lambda 2\omega$	$-\frac{n_1}{3}e_2+\frac{n_5}{2}e_5+\frac{n_6}{2}e_6$	$n_1e_1+n_5e_5+n_6e_6$
$\theta\lambda 2\omega\lambda 2$	$-\frac{n_1}{3}e_2-\frac{n_3}{3}e_4+\frac{n_5}{3}e_6$	$n_1e_1+n_3e_3-n_5e_5$
$\theta\lambda 2\omega\lambda 3$	$-\frac{n_1}{3}e_2+\frac{n_3}{2}e_3+\frac{n_4}{2}e_4$	$n_1e_1+n_3e_3+n_4e_4$
$\theta\lambda 2\omega\lambda 4$	$-\frac{n_1}{3}e_2-\frac{n_3}{3}e_4+a_5e_5+a_6e_6$	$n_1e_1-n_3e_3$
$\theta\lambda 2\omega\lambda 5$	$-\frac{n_1}{3}e_2$	n_1e_1
$\theta\lambda 3\omega$	$\frac{n_1}{2}e_1+\frac{n_2}{2}e_2+\frac{n_5}{3}e_6$	$n_1e_1+n_2e_2-n_5e_5$
$\theta\lambda 3\omega\lambda 2$	$\frac{n_1}{2}e_1+\frac{n_2}{2}e_2-\frac{n_3}{3}e_4$	$n_1e_1+n_2e_2+n_3e_3$
$\theta\lambda 3\omega\lambda 3$	$\frac{n_1}{2}e_1+\frac{n_2}{2}e_2+\frac{n_3}{2}e_3+\frac{n_4}{2}e_4+a_5e_5+a_6e_6$	$n_1e_1+n_2e_2+n_3e_3+n_4e_4$

Continued on next page

$\theta\lambda 3\omega\lambda 4$	$\frac{n_1}{2}e_1 + \frac{n_2}{2}e_2 - \frac{n_3}{3}e_4$	$n_1e_1 + n_2e_2 - n_3e_3$
$\theta\lambda 3\omega\lambda 5$	$\frac{n_1}{2}e_1 + \frac{n_2}{2}e_2 + \frac{n_5}{2}e_5 + \frac{n_6}{2}e_6$	$n_1e_1 + n_2e_2 + n_5e_5 + n_6e_6$
$\theta\lambda 4\omega$	$-\frac{n_1}{3}e_2$	$-n_1e_1$
$\theta\lambda 4\omega\lambda 2$	$-\frac{n_1}{3}e_2 - \frac{n_3}{3}e_4 + a_5e_5 + a_6e_6$	$-n_1e_1 + n_3e_3$
$\theta\lambda 4\omega\lambda 3$	$-\frac{n_1}{3}e_2 + \frac{n_3}{2}e_3 + \frac{n_4}{2}e_4$	$-n_1e_1 + n_3e_3 + n_4e_4$
$\theta\lambda 4\omega\lambda 4$	$-\frac{n_1}{3}e_2 - \frac{n_3}{3}e_4 + \frac{n_5}{3}e_6$	$-n_1e_1 - n_3e_3 + n_5e_5$
$\theta\lambda 4\omega\lambda 5$	$-\frac{n_1}{3}e_2 + \frac{n_5}{2}e_5 + \frac{n_6}{2}e_6$	$-n_1e_1 + n_5e_5 + n_6e_6$
$\theta\lambda 5\omega$	$a_5e_5 + a_6e_6$	0
$\theta\lambda 5\omega\lambda 2$	$-\frac{n_3}{3}e_4$	n_3e_3
$\theta\lambda 5\omega\lambda 3$	$\frac{n_3}{2}e_3 + \frac{n_4}{2}e_4 + \frac{n_5}{3}e_6$	$n_3e_3 + n_4e_4 + n_5e_5$
$\theta\lambda 5\omega\lambda 4$	$-\frac{n_3}{3}e_4 + \frac{n_5}{2}e_5 + \frac{n_6}{2}e_6$	$-n_3e_3 + n_5e_5 + n_6e_6$
$\theta\lambda 5\omega\lambda 5$	$\frac{n_5}{3}e_6$	$-n_5e_5$

The action of θ and ω on the fixed points (or tori) is given by

twisted sector	Action of θ	Action of ω
$\theta, \theta\lambda5$	I	I
$\theta\lambda2, \theta\lambda4$	$(n_1, n_5) \to (-n_1, -n_5)$	$n_5 \to -n_5$
$\theta\lambda3$	$(n_1, n_2, n_5, n_6) \to (n_1 + n_2, n_1, n_6, n_5 + n_6)$	$(n_5, n_6) \to (n_6, n_5 + n_6)$
$\omega, \omega\lambda5$	I	I
$\omega\lambda2, \omega\lambda4$	$n_5 \to -n_5$	$(n_3, n_5) \to (-n_3, -n_5)$
$\omega\lambda3$	$(n_5, n_6) \to (n_6, n_5 + n_6)$	$(n_3, n_4, n_5, n_6) \to (n_3 + n_4, n_3, n_6, n_5 + n_6)$
$\theta\omega, \theta\lambda5\omega\lambda5$	$n_5 \to -n_5$	$n_5 \to -n_5$
$\theta\omega\lambda2, \theta\lambda5\omega\lambda4$	$(n_5, n_6) \to (n_6, n_5 + n_6)$	$(n_3, n_5, n_6) \to (-n_3, n_6, n_5 + n_6)$
$\theta\omega\lambda3, \theta\lambda5\omega\lambda3$	$n_5 \to -n_5$	$(n_3, n_4, n_5) \to (n_3 + n_4, n_3, -n_5)$
$\theta\omega\lambda4, \theta\lambda5\omega\lambda2$	I	$n_3 \to -n_3$
$\theta\omega\lambda5, \theta\lambda5\omega$	I	I
$\theta\lambda2\omega, \theta\lambda4\omega\lambda5$	$(n_1, n_5, n_6) \to (-n_1, n_6, n_5 + n_6)$	$(n_5, n_6) \to\to (n_6, n_5 + n_6)$
$\theta\lambda2\omega\lambda2, \theta\lambda4\omega\lambda4$	$(n_1, n_5) \to (-n_1, -n_5)$	$(n_3, n_5) \to (-n_3, -n_5)$
$\theta\lambda2\omega\lambda3, \theta\lambda4\omega\lambda3$	$n_1 \to -n_1$	$(n_3, n_4) \to (n_3 + n_4, n_3)$
$\theta\lambda2\omega\lambda4, \theta\lambda4\omega\lambda2$	$n_1 \to -n_1$	$n_3 \to -n_3$
$\theta\lambda2\omega\lambda5, \theta\lambda4\omega$	$n_1 \to -n_1$	I
$\theta\lambda3\omega, \theta\lambda3\omega\lambda5$	$(n_1, n_2, n_5) \to (n_1 + n_2, n_1, -n_5)$	$n_5 \to -n_5$
$\theta\lambda3\omega\lambda2, \theta\lambda3\omega\lambda4$	$(n_1, n_2) \to (n_1 + n_2, n_1)$	$n_3 \to -n_3$
$\theta\lambda3\omega\lambda3$	$(n_1, n_2) \to (n_1 + n_2, n_1)$	$(n_3, n_4) \to (n_3 + n_4, n_3)$

The moduli dependent suppression factors for the non-zero Yukawa couplings amongst twisted sectors containing massless states are given in the conventions of table 4 by

Yukawa Coupling	λ_1	λ_2	λ_3
$T_{01}T_{14}T_{51}$	0	1 for $n_3(T_{14}) \neq 0$	0
$T_{01}T_{15}T_{50}$	0	0	0
$T_{01}T_{23}T_{42}$	0	$\mu(n_3, p_3, p_4)$	0
$T_{01}T_{24}T_{41}$	0	1 for $n_3(T_{24}) \neq 0$	0
$T_{01}T_{32}T_{33}$	0	$\mu(n_3, p_3, p_4)$	0
$T_{02}T_{13}T_{51}$	0	$\mu(n_3, p_3, p_4)$	0
$T_{02}T_{14}T_{50}$	0	0	1 for $n_5(T_{02}) \neq 0$
$T_{02}T_{22}T_{42}$	0	2 for $m_3(T_{02}) - n_3(T_{22}) \neq 0$	0
$T_{02}T_{23}T_{41}$	0	$\mu(n_3, p_3, p_4)$	1 for $n_5(T_{02}) \neq 0$
$T_{02}T_{24}T_{40}$	0	0	0
$T_{02}T_{31}T_{33}$	0	$\mu(n_3, p_3, p_4)$	0
$T_{02}T_{32}T_{32}$	0	2 for $m_3(T_{02}) - n_3(T_{32}) \neq 0$	1 for $n_5(T_{02}) \neq 0$
$T_{03}T_{12}T_{51}$	0	$\mu(n_3, p_3, p_4)$	0
$T_{03}T_{13}T_{50}$	0	0	$\mu(n_5, p_5, p_6)$
$T_{03}T_{21}T_{42}$	0	$\mu(n_3, p_3, p_4)$	0
$T_{03}T_{22}T_{41}$	0	$\mu(n_3, p_3, p_4)$	$\mu(n_5, p_5, p_6)$
$T_{03}T_{23}T_{40}$	0	0	$\mu(n_5, p_5, p_6)$
$T_{03}T_{30}T_{33}$	0	0	0
$T_{03}T_{31}T_{32}$	0	$\mu(n_3, p_3, p_4)$	$\mu(n_5, p_5, p_6)$
$T_{04}T_{11}T_{51}$	0	1 for $n_3(T_{04}) \neq 0$	0
$T_{04}T_{12}T_{50}$	0	0	$\mu(n_5, p_5, p_6)$
$T_{04}T_{20}T_{42}$	0	0	0
$T_{04}T_{21}T_{41}$	0	1 for $n_3(T_{04}) \neq 0$	$\mu(n_5, p_5, p_6)$
$T_{04}T_{22}T_{40}$	0	0	2 for $n_5(T_{04}) - n_5(T_{22}) \neq 0$

Continued on next page

$T_{04}T_{30}T_{32}$	0	0	$\mu(n_5, p_5, p_6)$
$T_{05}T_{10}T_{51}$	0	0	0
$T_{05}T_{11}T_{50}$	0	0	1 for $n_5(T_{11}) \neq 0$
$T_{05}T_{20}T_{41}$	0	0	1 for $n_5(T_{20}) \neq 0$
$T_{05}T_{21}T_{40}$	0	0	$\mu(n_5, p_5, p_6)$
$T_{05}T_{30}T_{31}$	0	0	$\mu(n_5, p_5, p_6)$
$T_{10}T_{14}T_{42}$	1 for $n_1(T_{42}) \neq 0$	0	0
$T_{10}T_{15}T_{41}$	1 for $n_1(T_{41}) \neq 0$	0	0
$T_{10}T_{23}T_{33}$	$\mu(n_1, p_1, p_2)$	0	0
$T_{10}T_{24}T_{32}$	$\mu(n_1, p_1, p_2)$	0	0
$T_{11}T_{13}T_{42}$	1 for $n_1(T_{42}) \neq 0$	$\mu(n_3, p_3, p_4)$	0
$T_{11}T_{14}T_{41}$	1 for $n_1(T_{41}) \neq 0$	1 for $n_3(T_{14}) \neq 0$	1 for $n_5(T_{11}) \neq 0$
$T_{11}T_{15}T_{40}$	1 for $n_1(T_{40}) \neq 0$	0	0
$T_{11}T_{22}T_{33}$	$\mu(n_1, p_1, p_2)$	$\mu(n_3, p_3, p_4)$	0
$T_{11}T_{23}T_{32}$	$\mu(n_1, p_1, p_2)$	$\mu(n_3, p_3, p_4)$	1 for $n_5(T_{11}) \neq 0$
$T_{11}T_{24}T_{31}$	$\mu(n_1, p_1, p_2)$	1 for $n_3(T_{24}) \neq 0$	0
$T_{12}T_{12}T_{42}$	1 for $n_1(T_{42}) \neq 0$	2 for $m_3(T_{12}) - n_3(T_{12}) \neq 0$	0
$T_{12}T_{13}T_{41}$	1 for $n_1(T_{41}) \neq 0$	$\mu(n_3, p_3, p_4)$	$\mu(n_5, p_5, p_6)$
$T_{12}T_{14}T_{40}$	1 for $n_1(T_{40}) \neq 0$	0	$\mu(n_5, p_5, p_6)$
$T_{12}T_{21}T_{33}$	$\mu(n_1, p_1, p_2)$	$\mu(n_3, p_3, p_4)$	0
$T_{12}T_{22}T_{32}$	$\mu(n_1, p_2, p_2)$	2 for $m_3(T_{12}) - n_3(T_{22}) \neq 0$	$\mu(n_5, p_5, p_6)$
$T_{12}T_{23}T_{31}$	$\mu(n_1, p_1, p_2)$	$\mu(n_3, p_3, p_4)$	$\mu(n_5, p_5, p_6)$
$T_{12}T_{24}T_{30}$	$\mu(n_1, p_1, p_2)$	0	0
$T_{13}T_{13}T_{40}$	1 for $n_1(T_{40}) \neq 0$	0	2 for $m_5(T_{13}) - n_5(T_{13}) \neq 0$
$T_{13}T_{20}T_{33}$	$\mu(n_1, p_1, p_2)$	0	0

Continued on next page

$T_{13}T_{21}T_{32}$	$\mu(n_1, p_1, p_2)$	$\mu(n_3, p_3, p_4)$	$\mu(n_5, p_5, p_6)$
$T_{13}T_{22}T_{31}$	$\mu(n_1, p_1, p_2)$	$\mu(n_3, p_3, p_4)$	2 for $m_5(T_{22}) - n_5(T_{13}) \neq 0$
$T_{13}T_{23}T_{30}$	$\mu(n_1, p_1, p_2)$	0	$\mu(n_5, p_5, p_6)$
$T_{14}T_{20}T_{32}$	$\mu(n_1, p_1, p_2)$	0	1 for $n_5(T_{20}) \neq 0$
$T_{14}T_{21}T_{31}$	$\mu(n_1, p_1, p_2)$	1 for $n_3(T_{14}) \neq 0$	$\mu(n_5, p_5, p_6)$
$T_{14}T_{22}T_{30}$	$\mu(n_1, p_1, p_2)$	0	$\mu(n_5, p_5, p_6)$
$T_{15}T_{20}T_{31}$	$\mu(n_1, p_1, p_2)$	0	0
$T_{15}T_{21}T_{30}$	$\mu(n_1, p_1, p_2)$	0	0
$T_{20}T_{22}T_{24}$	2 for $m_1(T_{20}) - n_1(T_{22}) \neq 0$	0	0
$T_{20}T_{23}T_{23}$	2 for $m_1(T_{20}) - n_1(T_{23}) \neq 0$	1 for $n_3(T_{20}) \neq 0$	0
$T_{21}T_{21}T_{24}$	2 for $m_1(T_{21}) - n_1(T_{21}) \neq 0$	1 for $n_3(T_{24}) \neq 0$	0
$T_{21}T_{22}T_{23}$	2 for $m_1(T_{21}) - n_1(T_{22}) \neq 0$	$\mu(n_3, p_3, p_4)$	$\mu(n_5, p_5, p_6)$
$T_{22}T_{22}T_{22}$	2 for $m_1(T_{22}) - n_1(T_{22}) \neq 0$	2 for $m_3(T_{22}) - n_3(T_{22}) \neq 0$	2 for $m_5(T_{22}) - n_5(T_{22}) \neq 0$